

Restore Deep Volume, Recover Attractive Faceline with Neuramis Deep/Volume and Coretox

Leaders Clinic, Seoul, Korea

Do Young Rhee, MD. PhD.

The aging process brings about various changes in the face, resulting in visible signs of aging. One prominent change involves the thinning and shrinking of subcutaneous fat, leading to a loss of volume and a sunken, gaunt appearance. This loss of fat can also accentuate the presence of wrinkles and fine lines. Furthermore, ptotic subcutaneous fat can accumulate in the jowl area, causing sagging skin and a less defined jawline. A youthful lower faceline is characterized by a smooth, well-defined jawline and neckline.

This presentation focuses on the characteristics of a youthful lower faceline and discusses methods for its restoration. Key features of a youthful faceline include a taut and defined jawline, a well-defined chin with a balanced profile, fullness in the cheek area, and a firm neck without sagging skin or wrinkles. Immediate restoration of facial volume can be achieved through the use of filler injections.

In addition to fillers, botulinum toxin injections play a crucial role in lower face contouring. One effective application of botulinum toxin injection is masseter reduction, which can help reshape the lower face and achieve a more desirable contour. By targeting the masseter muscles, which contribute to a square or bulky jawline, botulinum toxin injections can create a slimmer, more streamlined appearance.

The presentation specifically introduces the use of Neuramis Deep Lidocaine and Neuramis Volume Lidocaine for facial volume restoration through filler injections, along with Coretox, a botulinum toxin treatment, for masseter reduction and lower face contouring. Attendees will gain valuable insights into the benefits and techniques associated with these treatments, enabling them to provide comprehensive solutions for rejuvenating and enhancing the lower faceline.